




A service of the National Library of Medicine
and the National Institutes of Health



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Note: Performing your original search, *neointimal formation restenosis*, in PubMed will retrieve **471** citations.

Display Show Sort by Send to

All: 1 Review: 0

☐ 1: Heart Dis. 2002 Jan-Feb;4(1):18-25.

[Related Articles, Links](#)

Neointima formation after stent implantation in an experimental model of restenosis: polytetrafluoroethylene-covered versus uncovered stainless steel stents.

Sick PB, Brosteanu O, Niebauer J, Hehrlein C, Schuler G.

Heart Center, Department for Internal Medicine/Cardiology, University of Leipzig, Germany. sickp@medizin.uni-leipzig.de

The aim of the study was to assess whether stents covered with a membrane of polytetrafluoroethylene spanned over the meshes of a sandwich-configured double stent (n = 15) prevent migration of smooth muscle cells through stent spaces, leading to less neointima formation compared with uncovered stainless steel stents (n = 14) in iliac arteries of male Chinchilla Bastard rabbits (n = 18). Lumen stenosis was assessed by quantitative angiography immediately before the animals were killed 5 weeks after stent deployment. Neointima formation was quantified by histomorphometric analysis. There were large regional and individual differences in neointima formation, leading locally to a significantly higher degree of stenosis in covered stents (histologically, 76.0 +/- 13.7 vs. 62.9 +/- 12.9%; angiographically, 33.5 +/- 21.1 vs. 7.8 +/- 8.8%) compared with uncovered stents, though mean neointimal and lumen area values were not significantly different. In conclusion, polytetrafluoroethylene-covered stents do not prevent neointima formation compared with uncovered stents. Although the membrane reduces local smooth muscle cell migration, the neointima hyperplasia at the proximal and distal ends of a covered stent stimulates migration along its longitudinal axis. In this stent-restenosis model, regional and individual proliferation processes and not the membrane-covering strut-to-strut distances determine lumen restenosis.

PMID: 11975828 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)